arXiv:1403.4875v1 [math.RT] 19 Mar 2014

A THEOREM OF MŒGLIN-WALDSPURGER FOR COVERING GROUPS

SHIV PRAKASH PATEL

Abstract. Let E be a non-Archimedian local field of characteristic zero and residue characteristic p. Let $\mathbf{G}$ be a connected reductive group defined over E and π an irreducible admissible representation of $G = \mathbf{G}(E)$. A result of C. Mœglin and J.-L. Waldspurger (for $p \neq 2$) and S. Varma (for $p = 2$) states that the leading coefficient in the character expansion of π at the identity element of $\mathbf{G}(E)$ gives the dimension of a certain space of degenerate Whittaker forms. In this paper we generalize this result of Mœglin-Waldspurger to the setting of covering groups $\tilde{G}$ of G.

1. Introduction

Let E be a non-Archimedian local field of characteristic zero and $\mathbf{G}$ a connected split reductive group defined over E and $G = \mathbf{G}(E)$. Let $\mathfrak{g} = \mathrm{Lie}(\mathbf{G})$ be the Lie algebra of $\mathbf{G}$ and $\mathfrak{g} = \mathfrak{g}(E)$. Let (π, W) be an irreducible admissible representation of G. A theorem of F. Rodier, in [7], relates the dimension of the space of non-degenerate Whittaker functionals of π and coefficients in the character expansion of π around identity. More precisely, Rodier proves that if the residue characteristic of E is large enough and the group $\mathbf{G}$ is split then the dimension of any space of non-degenerate Whittaker functionals for (π, W) equals the coefficient in the character expansion of π at identity corresponding to an appropriate maximal nilpotent orbit in the Lie algebra $\mathfrak{g}$. Rodier proved his theorem assuming that the residue characteristic of E is large enough, in fact, greater than a constant which depends only on the root datum of $\mathbf{G}$. A theorem of C. Mœglin and J.-L. Waldspurger [4] generalizes this theorem of Rodier in several directions, in particular proving the theorem of Rodier for the fields E whose residue characteristic is odd and removing the assumption on $\mathbf{G}$ being split. The theorem of Mœglin-Waldspurger is a more precise statement about the coefficients appearing in the character expansion around identity and certain spaces of 'degenerate' Whittaker forms. In a recent work of S. Varma [8] this theorem has been proved for fields with even residue characteristic. So the theorem of Mœglin-Waldspurger is true for all connected reductive groups without any restriction on the residue characteristic of the field E. We now recall the theorem of Mœglin-Waldspurger. To state the theorem we need to introduce some notation. Let Y be a nilpotent element in $\mathfrak{g}$ and suppose

Date: March 20, 2014.
2010 *Mathematics Subject Classification.* Primary 22E50; Secondary 11F70, 11S37.
Key words and phrases. Covering groups, character expansion, degenerate Whittaker forms.

$\varphi : \mathbb{G}_m \longrightarrow \mathbf{G}$ is a one parameter subgroup satisfying

$$Ad(\varphi(t))Y = t^{-2}Y. \tag{1}$$

Associated to such a pair (Y, φ) one can define a certain space $\mathcal{W}_{(Y,\varphi)}$, called the space of degenerate Whittaker forms of (π, W) relative to (Y, φ) (see Section 4 for the definition).

Define $\mathcal{N}_{Wh}(\pi)$ to be the set of nilpotent orbits $\mathcal{O}$ of $\mathfrak{g}$ for which there exists an element $Y \in \mathcal{O}$ and a φ satisfying (1) such that the space $\mathcal{W}_{(Y,\varphi)}$ of degenerate Whittaker forms relative to the pair (Y, φ) is non-zero.

Recall that the character expansion of (π, W) around identity is a sum $\sum_{\mathcal{O}} c_{\mathcal{O}} \widehat{\mu_{\mathcal{O}}}$, where $\mathcal{O}$ varies over the set of nilpotent orbits of $\mathfrak{g}$, $c_{\mathcal{O}} \in \mathbb{C}$ and $\widehat{\mu_{\mathcal{O}}}$ is the Fourier transform of a suitably chosen measure $\mu_{\mathcal{O}}$ on $\mathcal{O}$. One defines $\mathcal{N}_{tr}(\pi)$ to be the set of nilpotent orbits $\mathcal{O}$ of $\mathfrak{g}$ such that the corresponding coefficient $c_{\mathcal{O}}$ in the character expansion of π around identity is non zero.

We have the standard partial order on the set of nilpotent orbits in $\mathfrak{g}$: $\mathcal{O}_1 \leq \mathcal{O}_2$ if $\mathcal{O}_1 \subset \overline{\mathcal{O}_2}$. Let $\mathrm{Max}(\mathcal{N}_{Wh}(\pi))$ and $\mathrm{Max}(\mathcal{N}_{tr}(\pi))$ denote the set of maximal element in $\mathcal{N}_{Wh}(\pi)$ and $\mathcal{N}_{tr}(\pi)$ respectively with respect to this partial order. Then the main theorem of Mœglin-Waldspurger in Chapter I of [4] is as follows:

Theorem 1. *Let* $\mathbf{G}$ *be a connected reductive group defined over* E*. Let* π *be an irreducible admissible representation of* $G = \mathbf{G}(E)$ *then*

$$\mathrm{Max}(\mathcal{N}_{Wh}(\pi)) = \mathrm{Max}(\mathcal{N}_{tr}(\pi)).$$

Moreover, if $\mathcal{O}$ *is an element in either of these sets, then for any* (Y, φ) *as above with* $Y \in \mathcal{O}$ *we have*

$$c_{\mathcal{O}} = \dim \mathcal{W}_{(Y,\varphi)}.$$

If one considers the case of the pair (Y, φ) with Y a 'regular' nilpotent element then the above theorem of Mœglin-Waldspurger specializes to Rodier's theorem.

In this paper we generalize the theorem of Mœglin-Waldspurger to the setting of a covering group $\tilde{G}$ of G. Let μ_r be the group of r-th roots of unity in $\mathbb{C}^\times$. An r-fold covering group $\tilde{G}$ of G is a central extension of G by μ_r given by the following short exact sequence

$$1 \longrightarrow \mu_r \longrightarrow \tilde{G} \longrightarrow G \longrightarrow 1. \tag{2}$$

The representations of $\tilde{G}$ on which μ_r acts by the natural embedding $\mu_r \hookrightarrow \mathbb{C}^\times$ are called genuine representations. The definition of the space of degenerate Whittaker forms of a representation of G involves only unipotent groups. Since the covering $\tilde{G} \longrightarrow G$ splits over any unipotent subgroup of G in a unique way, see [5], this makes it possible to define the space of degenerate Whittaker forms for any genuine smooth representation (π, W) of $\tilde{G}$.

In particular, it makes sense to talk of the set $\mathcal{N}_{Wh}(\pi)$.

The existence of character expansion of an admissible genuine representation (π, W) of $\tilde{G}$ has been proved by Wen-Wei Li in [1]. At identity, the Harish-Chandra-Howe character expansion of an irreducible genuine representation has the same form and therefore we have $\mathcal{N}_{\text{tr}}(\pi)$. This makes it possible to have an analogue of Theorem 1 to the setting of covering groups. The main aim of this paper is to prove the following.

Theorem 2. *Let π be an irreducible admissible genuine representation of $\tilde{G}$. Then*

$$\text{Max}(\mathcal{N}_{Wh}(\pi)) = \text{Max}(\mathcal{N}_{tr}(\pi)).$$

Moreover, if $\mathcal{O}$ is an element in either of these sets, then for any (Y, φ) as above with $Y \in \mathcal{O}$ we have

$$c_{\mathcal{O}} = \dim \mathcal{W}_{(Y,\varphi)}.$$

We will use the work of Mœglin-Waldspurger [4] and to accommodate the even residue characteristic case, we follow Varma [8]. Let us describe some ideas involved in the proof. Let Y be a nilpotent element in $\mathfrak{g}$ and φ a one parameter subgroup as above. Let $\mathfrak{g}_i$ be the eigenspace of weight i under the action of $\mathbb{G}_m$ on $\mathfrak{g}$ via $\text{Ad} \circ \varphi$. One can attach a parabolic subgroup $\mathbf{P} := \oplus_{i \geq 0} \mathfrak{g}_i$ of $\mathbf{G}$ with unipotent radical $\mathbf{N} = \oplus_{i>0} \mathfrak{g}_i$. The one parameter subgroup φ also determines a parabolic subgroup $\mathbf{P}^- = \oplus_{i \leq 0} \mathfrak{g}_i$ opposite to $\mathbf{P}$. For simplicity, assume $\mathfrak{g}_1 = 0$ for the purpose of the introduction. Then $\mathbf{N} = \oplus_{i \geq 2} \mathfrak{g}_i$ and $\chi : \gamma \mapsto \psi(B(Y, \log \gamma))$ defines a character of $N = \mathbf{N}(E)$, where B is an $\text{Ad}(G)$-invariant non-degenerate symmetric bilinear form on $\mathfrak{g}$ and ψ is an additive character of E. In this case (i.e., $\mathfrak{g}_1 = 0$), the space of degenerate Whittaker forms $\mathcal{W}_{(Y,\varphi)}$ is defined to be the twisted Jacquet module of π with respect to (N, χ). In the case where $\mathfrak{g}_1 \neq 0$, the definition of $\mathcal{W}_{(Y,\varphi)}$ needs to be appropriately modified (see Section 4).

On the other hand, to the pair (Y, φ) one attaches certain open compact subgroups G_n of G for large n and certain characters χ_n of G_n. One then proves that the covering $\tilde{G} \longrightarrow G$ splits over G_n for large n, so that G_n can be seen as subgroups of $\tilde{G}$ as well. Let $t := \varphi(\varpi)$ and $\tilde{t}$ be any lift of t in $\tilde{G}$. It turns out that $\tilde{t}^{-n} G_n \tilde{t}^n \cap N$ becomes "arbitrarily large" subgroup of N and $\tilde{t}^{-n} G_n \tilde{t}^n \cap P^-$ gets "arbitrarily small" subgroup of P^-, as n becomes large. For large n, the characters χ_n have been so defined that that the character $\chi_n' := \chi_n \circ \text{Int}(\tilde{t}^n)$ restricted to $\tilde{t}^{-n} G_n \tilde{t}^n \cap N$ agrees with χ. Using Harish-Chandra-Howe character expansion one proves that the dimension of (G_n, χ_n)-isotypic component of W is equal to $c_{\mathcal{O}}$ for large enough n, where $\mathcal{O}$ is the nilpotent orbit of Y in $\mathfrak{g}$. Note that the (G_n, χ_n)-isotypic component of W and the $(\tilde{t}^{-n} G_n \tilde{t}^n, \chi_n \circ \text{Int}(\tilde{t}^n))$-isotypic component of W are isomorphic as vector spaces. Finally one proves that there is a natural isomorphism between $(\tilde{t}^{-n} G_n \tilde{t}^n, \chi_n \circ \text{Int}(\tilde{t}^n))$-isotypic component of W and $\mathcal{W}_{(Y,\varphi)}$.

Although the methods used in the paper are not new and heavily depend on the proofs in the linear case, the result is useful in the study of representation theory of covering groups. Author himself has made use of this result in his thesis, where he attempts to generalize a

result of D. Prasad in [6] in the setting of covering groups, namely, in the harmonic analysis relating the pairs $(\widetilde{\mathrm{GL}_2(E)}, \mathrm{GL}_2(F))$ and $(\widetilde{\mathrm{GL}_2(E)}, D_F^\times)$, where E/F is a quadratic extension of non-Archimedian local field, D_F is the quaternion division algebra with center F and $\widetilde{\mathrm{GL}_2(E)}$ is a certain two fold cover of $\mathrm{GL}_2(E)$.

Let us briefly give an outline of the organization of the paper. In Section 2, we recall the definition of the subgroups G_n and state some properties of the character χ_n. In Section 3, we recall splitting of the covering groups over G_n and describe an appropriate choice of the splitting over the subgroup G_n for large n. In Section 4 we give the definition of the space of degenerate Whittaker forms and describe important set up to prove the main theorem. In Section 5, we transfer some results from linear groups to covering groups in a few lemmas and based on these lemmas we prove the main theorem.

Acknowledgements: Author would like to express his gratitude to Professor D. Prasad and Professor Sandeep Varma for their numerous help and suggestions at various points. Without their help and continuous encouragement this paper would not have been possible.

2. Subgroups G_n and characters χ_n

In this section, we recall a certain sequence of subgroups G_n of G, which form a basis of neighbourhoods at identity and certain characters $\chi_n : G_n \longrightarrow \mathbb{C}^\times$. Although the objects involved in this section were defined for linear groups in [4, 8], we will lift them to covering groups in a suitable way in Section 3 and work with these lifts in this paper.

Let $\mathfrak{O}_E$ denotes the ring of integers in E. We fix an additive character ψ of E with conductor $\mathfrak{O}_E$. Fix an $\mathrm{Ad}(G)$-invariant non-degenerate symmetric bilinear form $B : \mathfrak{g}\times\mathfrak{g} \longrightarrow E$.

Let Y be a nilpotent element in $\mathfrak{g}$. Choose a one parameter subgroup $\varphi : \mathbb{G}_m \longrightarrow \mathbf{G}$ satisfying

$$\mathrm{Ad}(\varphi(s))Y = s^{-2}Y, \forall s \in \mathbb{G}_m. \tag{3}$$

Existence of such a φ is known from the theory of $\mathfrak{sl}_2$-triplets. But there are examples which do not come from this theory. We note that for a given nilpotent element $Y \in \mathfrak{g}$ the existence of φ is guaranteed by the theory of $\mathfrak{sl}_2$-triplets.
For $i \in \mathbb{Z}$, define

$$\mathfrak{g}_i = \{X \in \mathfrak{g} : \mathrm{Ad}(\varphi(s))X = s^i X, \forall s \in \mathbb{G}_m\}.$$

Set

$$\mathfrak{n} := \mathfrak{n}^+ := \oplus_{i>0}\mathfrak{g}_i, \mathfrak{n}^- := \oplus_{i<0}\mathfrak{g}_i, \mathfrak{p}^- := \oplus_{i\leq 0}\mathfrak{g}_i.$$

The parabolic subgroup $\mathbf{P}^-$ of $\mathbf{G}$ stabilizing $\mathfrak{n}^-$ has $\mathfrak{p}^-$ as its Lie algebra. Let $\mathbf{N} = \mathbf{N}^+$ be the unipotent subgroup of $\mathbf{G}$ having the Lie algebra $\mathfrak{n}$.

Let $G(Y)$ be the centralizer of Y in G and $Y^\#$ the centralizer of Y in $\mathfrak{g}$. The G-orbit $\mathcal{O}_Y$ of Y can be identified with $G/G(Y)$ and therefore its tangent space at Y can be identified with

$\mathfrak{g}/Y^{\#}$. Note that

$$\begin{aligned} Y^{\#} &= \{X \in \mathfrak{g} : [X, Y] = 0\} \\ &= \{X \in \mathfrak{g} : B([X, Y], Z) = 0, \forall Z \in \mathfrak{g}\} \\ &= \{X \in \mathfrak{g} : B(Y, [X, Z]) = 0, \forall Z \in \mathfrak{g}\}. \end{aligned}$$

The bilinear form B induces a non-degenerate alternating form $B_Y : \mathfrak{g}/Y^{\#} \times \mathfrak{g}/Y^{\#} \longrightarrow E$ defined by $B_Y(X_1, X_2) = B(Y, [X_1, X_2])$.

Let $L \subset \mathfrak{g}$ be a lattice satisfying the following conditions:

(1) $[L, L] \subset L$,
(2) $L = \oplus_{i\in\mathbb{Z}} L_i$, where $L_i = L \cap \mathfrak{g}_i$,
(3) The lattice L/L_Y, where $L_Y = L \cap Y^{\#}$, is self dual (i.e. $(L/L_Y)^{\perp} = L/L_Y$) with respect to B_Y. (For any vector space V with a non-degenerate bilinear form B and a lattice M in V, $M^{\perp} := \{X \in V : B(X, Y) \in \mathfrak{O}_E, \forall Y \in V\}$.)

A lattice L satisfying the above properties can be chosen by taking a suitable basis of all $\mathfrak{g}_i$'s, see [4]. Now we summarize a few well known properties of the exponential map, and use them to define subgroups G_n and their Iwahori decompositions.

Lemma 1. (1) *There exists a positive integer A such that* exp *is defined and injective on $\varpi^A L$, with inverse* log.
(2) *The image $G_n := \exp(\varpi^n L)$ is a subgroup of G for all $n \geq A$.*
(3) *Set $P_n^- = \exp(\varpi^n L \cap \mathfrak{p}^-)$ and $N_n = \exp(\varpi^n L \cap \mathfrak{n})$. Then we have an Iwahori factorization*

$$G_n = P_n^- N_n.$$

We will be working with a certain character χ_n of G_n, which we recall in the next lemma.

Lemma 2. *For large values of n there exists a character χ_n of G_n, whose restriction to $\exp((Y^{\#} \cap \varpi^n L) + \varpi^{n+\mathrm{val}2} L)$ coincides with $\gamma \mapsto \psi(B(\varpi^{-2n} Y, \log \gamma))$. If P_n^- is as in Lemma 1, the character χ_n can be chosen so that*

$$\chi_n(p) = 1, \forall p \in P_n^-.$$

For a proof of this lemma and other properties of this character χ_n see Lemma 5 in [8].

Remark 1. If $p \neq 2$, then the map $\gamma \mapsto \psi(B(\varpi^{-2n} Y, \log \gamma))$ itself defines a character of G_n for large n and satisfies the properties stated in Lemma 2. For $p \neq 2$, clearly there is just one character χ_n of G_n with these properties. But when $p = 2$, the number of such characters is greater than one, for more details see [8].

3. Covering groups

Let μ_r be the group of r-th roots of unity in $\mathbb{C}$. Consider an r-fold covering $\tilde{G}$ of G which is a central extension of the group G by μ_r:

$$1 \longrightarrow \mu_r \longrightarrow \tilde{G} \longrightarrow G \longrightarrow 1.$$

Lemma 3. (1) *The covering $\tilde{G} \longrightarrow G$ splits uniquely over any unipotent subgroup of G.*

(2) *For large enough n the covering $\tilde{G} \longrightarrow G$ splits over G_n. Moreover, there is a splitting s of $\tilde{G} \longrightarrow G$ restricted to $\cup_{g\in G} gG_ng^{-1}$ such that $s(hth^{-1}) = hs(t)h^{-1}$ for all $h \in G$.*

Proof. (1) This is well known, see [5]. For a simpler proof, in the case when E has characteristic zero, see Section 2.2 of [2].

(2) Recall that the subgroups G_n form a basis of neighbourhoods of identity. It is well known that the covering $\tilde{G} \longrightarrow G$ splits over a neighbourhood of identity. Therefore for large enough n, the covering splits over G_n. There are more than on possible splitting of the cover $\tilde{G} \longrightarrow G$ over G_n. If a splitting is fixed, then any other splitting over G_n will differ from the above splitting by a character $G_n \longrightarrow \mu_r$.
Fix some m such that the covering splits over $G_m = \exp(\varpi^m L)$. As mentioned above, any two splitting over the subgroup G_m will differ by a character $G_m \longrightarrow \mu_r$ and any such character is trivial over

$$G_m^r := \{g^r : g \in G_m\}.$$

Hence all the possible splittings over G_m agree on G_m^r. The subset G_m^r is a subgroup of G_m as it equals $\exp(r \cdot \varpi^m L)$. Let $g, h \in G$. Then

$$(gG_mg^{-1} \cap hG_mh^{-1}) \supset (gG_m^rg^{-1} \cap hG_m^rh^{-1}).$$

This implies that any two splittings of $\tilde{G} \longrightarrow G$ restricted to $gG_m^rg^{-1} \cap hG_m^rh^{-1}$ one coming from the restriction of a splitting of $\tilde{G} \longrightarrow G$ over gG_mg^{-1} and the other coming from the restriction of a splitting over hG_mh^{-1} are the same.. Now choose A' so large such that $G_n \subset G_m^r$ for $n \geq A'$. We fix the splitting of G_n which comes from that of the restriction of G_m^r. This gives us a splitting over $\cup_{g\in G} gG_ng^{-1}$. □

Using this splitting we get that exponential map is defined from a small enough neighbourhood of $\mathfrak{g}$ to $\tilde{G}$, namely the usual exponential map composed with this splitting, which one can use to define the character expansion of an irreducible admissible genuine representation (π, W) of $\tilde{G}$, which has been done by Wen-Wei Li in [1].

Remark 2. If r is co-prime to p, then as G_n is a pro-p group and $(r, p) = 1$, there is no non-trivial character from G_n to μ_r. In that situation, the splitting in the above lemma is unique.

From now onwards, for large enough n, we treat G_n not only as a subgroup of G but also as one of $\tilde{G}$, with the above specified splitting. In other words, for the covering group $\tilde{G}$ (as in the linear case) we have a sequence of pairs (G_n, χ_n) using the splitting specified above which satisfies the properties described in Section 2.

Definition 1. Let $H \subset G$ be an open subgroup and $s : H \hookrightarrow \tilde{G}$ be a splitting. Then for any $\phi \in C_c^\infty(G)$ with $\mathrm{supp}(\phi) \subset H$ define $\tilde{\phi}_s \in C_c^\infty(\tilde{G})$ as follows:

$$\tilde{\phi}_s(g) := \begin{cases} \phi(g), & \text{if } g \in s(H) \\ 0, & \text{if } g \in \tilde{G} \backslash s(H) \end{cases}$$

Note that this definition depends upon the choice of splitting. Whenever the splitting is clear in the context or it has been fixed and there is no confusion we write just $\tilde{\phi}$ instead of $\tilde{\phi}_s$ and H for $s(H)$. Recall that the convolution $\phi * \phi'$ for $\phi, \phi' \in C_c^\infty(G)$ is defined by

$$\phi * \phi'(x) = \int_G \phi(xy^{-1})\phi'(y)\, dy.$$

Observe that

$$\mathrm{supp}(\phi * \phi') \subset \mathrm{supp}(\phi) \cdot \mathrm{supp}(\phi'),$$

which implies the lemma below.

Lemma 4. *Let H be an open subgroup of G such that the covering $\tilde{G} \to G$ has a splitting over H, say, $s : H \hookrightarrow \tilde{G}$, satisfying $s(xy) = s(x)s(y)$ whenever x, y are in H. If $\phi, \phi' \in C_c^\infty(G)$ are such that supp(ϕ) and supp(ϕ') are contained in H, then we have*

$$\widetilde{\phi * \phi'} = \tilde{\phi} * \tilde{\phi}'.$$

4. Degenerate Whittaker forms

In this section we give the definition of degenerate Whittaker forms for a smooth genuine representation π of $\tilde{G}$. This is an adaptation of Section I.7 of [4] and Section 5 of [8].

Recall that $N^2 = \exp(\oplus_{i \geq 2} \mathfrak{g}_i)$ and $N' = \exp(\mathfrak{g}_1 \cap Y^\#)N^2$ are normal subgroups of N. Let H be the Heisenberg group defined with $\mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#) \times E$ as underlying set using the symplectic from induced by B_Y. Consider the map $N \longrightarrow H$ given by

$$\exp(X) \mapsto (\overline{X}, B(Y, X)),$$

where $\overline{X}$ is the image of the $\mathfrak{g}_1$ component of X in $\mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#)$. The Campbell-Hausdorff formula implies that the above map is a homomorphism with the following kernel

$$N'' = \{n \in N' : B(Y, \log n) = 0\}.$$

Let $\chi : N' \longrightarrow \mathbb{C}^\times$ be defined by $\gamma \mapsto \psi \circ B(Y, \log \gamma)$. Note that $\gamma \mapsto B(Y, \log \gamma) \in E \cong \{0\} \times E \subset H$ induces an isomorphism $N'/N'' \cong E$.

We note that the cover $\tilde{G} \longrightarrow G$ splits uniquely over the subgroups N, N' and N''. We denote the images of these splittings inside $\tilde{G}$ by the same letters. For a smooth genuine representation (π, W) of $\tilde{G}$ we define

$$N_\chi^2 W = \{\pi(n)w - \chi(n)w : w \in W, n \in N^2\}$$

and

$$N'_\chi W = \{\pi(n)w - \chi(n)w : w \in W, n \in N'\}.$$

Note that N normalizes χ, therefore $H = N/N''$ acts on $W/N'_\chi W$ in a natural way. This action restricts to N'/N'' (the center of N/N'') as multiplication by the character χ. Let $\mathcal{S}$ be the unique irreducible representation of the Heisenberg group H with central character χ.

Definition 2. Define the space of degenerate Whittaker forms for (π, W) associated to (Y, φ) to be

$$\mathcal{W} := \mathrm{Hom}_H(\mathcal{S}, W/N'_\chi W).$$

Remark 3. If $\mathfrak{g}_1 = 0$, then $N = N' = N^2$. In this case, $\mathcal{W} \cong W/N_\chi W$ is the (N, χ)-twisted Jacquet functor.

Definition 3. For a smooth representation (π, W) of $\tilde{G}$ define $\mathcal{N}_{\mathrm{Wh}}(\pi)$ to be the set of nilpotent orbits $\mathcal{O}$ of $\mathfrak{g}$ such that there exists $Y \in \mathcal{O}$ and φ as in Equation 3, such that the space of degenerate Whittaker forms for π associated to (Y, φ) is non-zero.

As $\mathfrak{g}_1$ is a symplectic vector space and L/L_Y is self dual, it follows that $L_H := (L \cap \mathfrak{g}_1)/(L \cap \mathfrak{g}_1 \cap Y^\#)$ is a self dual lattice in the symplectic vector space $H/Z(H) \cong \mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#)$.

Recall the definition of the Heisenberg group H: for $X, Z \in \mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#)$ and $a, b \in E$,

$$(X, a)(Z, b) = (X + Z, a + b + \frac{1}{2}B_Y(X, Z)).$$

As ψ is trivial on $\mathfrak{O}_E$, it follows that one can extend the character ψ of $E \cong Z(H)$ to a character of the inverse image of $2L_H$ under $H \longrightarrow \mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#)$ by defining it to be trivial on $2L_H \times \{0\} \subset H$. From Lemma 4 in [8], this character can be extended to a character $\tilde{\chi}$ on the inverse image H_0 of L_H under the natural map $H \longrightarrow \mathfrak{g}_1/(\mathfrak{g}_1 \cap Y^\#)$.

Remark 4. There are one parameter subgroups φ which do not arise from $\mathfrak{sl}_2$-triplets. If φ arises from $\mathfrak{sl}_2$-triplets, then it is easy to see that $Y^\# \subset \oplus_{i \leq 0}\mathfrak{g}_i$. In particular we have $\mathfrak{g}_1 \cap Y^\# = \{0\}$ and hence the Heisenberg group $H = \mathfrak{g}_1 \times E$.

Then, by Section I.3 of [3], one knows that $\mathcal{S} = \mathrm{ind}_{H_0}^H \tilde{\chi}$, induction with compact support. Since H_0 is an open subgroup of the locally profinite group H, we have the following form of the Frobenius reciprocity law:

$$\mathrm{Hom}_H(\mathcal{S}, \tau) = \mathrm{Hom}_H(\mathrm{ind}_{H_0}^H \tilde{\chi}, \tau) = \mathrm{Hom}_{H_0}(\tilde{\chi}, \tau\mid_{H_0})$$

for any smooth representation τ of H. Thus, in the category of representations of N on which N' acts via the character χ, the functor $\mathrm{Hom}_H(\mathcal{S}, -)$ amounts to taking the $\tilde{\chi}\mid_{H_0}$-isotypic component. Since H_0 is compact modulo the center, this functor is exact. Thus we have

$$\mathcal{W} = \mathrm{Hom}_H(\mathcal{S}, W/N'_\chi W) \cong (W/N'_\chi W)^{(H_0, \tilde{\chi})}.$$

where $(W/N'_\chi W)^{(H_0,\tilde{\chi})}$ denotes the $(H_0,\tilde{\chi})$-isotypic component of $W/N'_\chi W$.

Recall that we have defined certain characters χ_n's in Section 2 and now we have a character $\tilde{\chi}$. We need to choose them in a compatible way. First we fix a character $\tilde{\chi}$ and consider it as a character of $\exp(\mathfrak{g}_1 \cap L)N'$ in the obvious way (as $\exp(\mathfrak{g}_1 \cap L)N'$ is the inverse image of H_0 under $N \longrightarrow H$). Let $t := \varphi(\varpi) \in G$. Let $\tilde{t} \in \tilde{G}$ be any lift of t in $\tilde{G}$. Let

$$G'_n = \mathrm{Int}(\tilde{t}^{-n})(G_n), P'_n = \mathrm{Int}(\tilde{t}^{-n})(P_n^-) \text{ and } V'_n = \mathrm{Int}(\tilde{t}^{-n})(N_n).$$

It can be easily verified that V'_n contains $\exp(\mathfrak{g}_1 \cap L)$. We also have $V'_n \subset V'_m$ for large m, n with $n \leq m$. Moreover

$$\exp(\mathfrak{g}_1 \cap L)N^2 = \bigcup_{n\geq 0} V'_n.$$

It can also be verified easily that $\tilde{\chi} \circ \mathrm{Int}(\tilde{t}^{-n})$ restricts to a character of N_n that extends the character on $N_{n+\mathrm{val}2}N'_n$ given by $\gamma \mapsto \psi(B(\varpi^{-2n}Y, \log\gamma))$. Now define

$$\chi_n(pv) = \tilde{\chi}(\tilde{t}^{-n}v\tilde{t}^n), \forall p \in P_n^- \text{ and } \forall v \in V'_n. \tag{4}$$

Lemma 5 (Lemma 6 in [8])**.** *Let χ_n be as defined in Equation 4. Then χ_n is a character of G_n and satisfies the properties stated in Lemma 2.*

Define a character χ'_n on G'_n as follows:

$$\chi'_n := \chi_n \circ \mathrm{Int}(\tilde{t}^n).$$

Remark 5. The characters χ_n have been so defined that χ'_n agree with χ on the intersection of their domains, namely, for large n we have,

$$\chi'_n \mid_{V'_n} = \tilde{\chi} \mid_{V'_n} .$$

In particular, $\chi'_n \mid_{\exp(L\cap\mathfrak{g}_1)} = \tilde{\chi} \mid_{\exp(L\cap\mathfrak{g}_1)}$. One can also see that χ'_n and χ'_m (for large n, m) agree on $G'_n \cap G'_m$, because they agree on $V'_n \cap V'_m$ and also on $P'_n \cap P'_m$ (being trivial on it).

Set

$$W_n := \{w \in W \mid \pi(\gamma)w = \chi_n(\gamma)w, \forall \gamma \in G_n\} \tag{5}$$

and

$$W'_n := \{w \in W \mid \pi(\gamma)w = \chi'_n(\gamma)w, \forall \gamma \in G'_n\} = \pi(\tilde{t}^{-n})W_n \tag{6}$$

For large m, n define the map $I'_{n,m} : W'_n \longrightarrow W'_m$ by

$$I'_{n,m}(w) = \int_{G'_m} \chi'_m(\gamma^{-1})\pi(\gamma)w\, d\gamma. \tag{7}$$

Let m, n be large with $m > n$. Since χ'_n is trivial on $P'_n \supset P'_m$ and since $G'_m = P'_m V'_m$ and for a convenient choice of measures we have

$$\begin{aligned} I'_{n,m}(w) &= \int_{V'_m} \chi'_m(x^{-1})\pi(x)w\, dx \\ &= \int_{\exp(\mathfrak{g}_1\cap L)} \tilde{\chi}^{-1}(\exp X)\pi(\exp X) \int_{N^2\cap G'_m} \chi(x^{-1})\pi(x)w\, dx\, dX. \end{aligned}$$

Now using the fact that $\exp(\mathfrak{g}_1 \cap L)$ lies in G'_n for large n and that it normalizes the character $\chi|_{N^2}$, we get

$$\begin{aligned} I'_{n,m}(w) &= \int_{N^2 \cap G'_m} \chi(x^{-1})\pi(x)w\, dx \\ &= \int_{N' \cap G'_m} \chi(x^{-1})\pi(x)w\, dx. \end{aligned}$$

From this the following is clear for large n, m with $m > n$

$$I'_{n,m} = I'_{n+1,m} \circ I'_{n,n+1}. \tag{8}$$

For large n, the above equation gives that $\ker I'_{n,m} \subset \ker I'_{n,p}$ for $n < m \leq p$. Set $W'_{n,\chi} := \cup_{m>n} \ker I'_{n,m}$. We get natural maps $j_n : W'_n/W'_{n,\chi} \longrightarrow W/N^2_\chi W$ and $j'_n : W'_n/W'_{n,\chi} \longrightarrow W/N'_\chi W$ that gives the following diagram:

$$\begin{array}{ccc} W'_n/W'_{n,\chi} & \xrightarrow{\quad j'_n \quad} & W/N'_\chi W \\ & \searrow^{j_n} \quad \nearrow_{\exists \text{ natural}} & \\ & W/N^2_\chi W & \end{array} \tag{9}$$

By the compatibility between χ'_n and $\tilde{\chi}$, it is easy to see that the image of j'_n is contained in $(W/N'_\chi W)^{(H_0,\tilde{\chi})}$. Let $w \in W$ such that the image $\bar{w}$ of w in $W/N'_\chi W$ belongs to $(W/N'_\chi W)^{(H_0,\tilde{\chi})}$. For large n, P'_n acts trivially on w, as (π, W) is smooth. Since $G'_n = P'_n V'_n = V'_n P'_n$, the element

$$\int_{V'_n} \chi'_n(x^{-1})\pi(x)w\, dx$$

belongs to W'_n. As χ'_n and χ are compatible, it can be seen that its image in $W/N'_\chi W$ is $\bar{w}$. This gives us the following lemma.

Lemma 6. *Let (Y, φ) be arbitrary. Then any element of $(W/N'_\chi W)^{(H_0,\chi)}$ belongs to $j'_n(W'_n)$ for all sufficiently large n. In particular, if $\mathcal{W} \neq 0$ then, for large n, W_n and W'_n are non-zero.*

5. Main theorem

Now recall that, by the work of Wen-Wei Li [1], the Harish-Chandra-Howe character expansion of an irreducible admissible genuine representation of $\tilde{G}$ at the identity element has an expression of the same form as that of an irreducible admissible representation of a linear group. The proof of the following lemma for a covering group follows verbatim that of Proposition I.11 in [4] and Proposition 1 in [8].

Proposition 1. *Let $\mathcal{W}$ be the space of degenerate Whittaker forms for π with respect to a given (Y, φ). If $\mathcal{W} \neq 0$ then there exists a nilpotent orbit $\mathcal{O}$ in $\mathcal{N}_{\mathrm{tr}}(\pi)$ such that $\mathcal{O}_Y \leq \mathcal{O}$ (i.e., $Y \in \bar{\mathcal{O}}$).*

Let the function $\phi_n : G \longrightarrow \mathbb{C}$ be defined by

$$\phi_n(\gamma) = \begin{cases} \chi_n(\gamma^{-1}), & \text{if } \gamma \in G_n \\ 0, & \text{otherwise.} \end{cases}$$

Consider the corresponding function $\tilde{\phi}_n : \tilde{G} \longrightarrow \mathbb{C}$. Write the character expansion at the identity element as follows:

$$\Theta_\pi = \sum_O c_O \widehat{\mu_O}.$$

Choose n large enough so that the above expansion is valid over G_n and then evaluate Θ_π at the function $\tilde{\phi}_n$. As $\pi(\tilde{\phi}_n)$ is a projection from W to W_n, by definition we get $\Theta_\pi(\tilde{\phi}_n) = \operatorname{trace} \pi(\tilde{\phi}_n) = \dim W_n$. Now assume that (Y, φ) is such that O_Y is a maximal element in $\mathcal{N}_{tr}(\pi)$. On the other hand, if we evaluate $\sum_O c_O \widehat{\mu_O}(\tilde{\phi}_n)$, it turns out that $\widehat{\mu_O}(\tilde{\phi}_n)$ is zero unless $O = O_Y$. In addition, if we fix a G-invariant measure on O_Y as in I.8 of [4] (for more details about this invariant measure see Section 3 of [8]), we get the following lemma.

Lemma 7. *[Lemma I.12 in* [4] *and Lemma 7 in* [8]*]*
If (Y, φ) is such that O_Y is a maximal element of $\mathcal{N}_{\mathrm{tr}}(\pi)$. Then for large n,

$$\dim W_n = c_{O_Y}.$$

In particular, the dimension of W_n is finite and independent of n, for large n.

From Lemma 6 we know that every vector in $\mathcal{W}$ is in the image of j'_n for large n. In particular, if W_n is finite dimensional, we get that the map j'_n is surjective. Moreover, we have the following lemma whose proof is verbatim that of Corollary I.14 in [4] and Lemma 8 in [8] in the case of a linear group.

Lemma 8. *Let (Y, φ) is such that O_Y is a maximal element of $\mathcal{N}_{\mathrm{tr}}(\pi)$. Then for large n, the maps j_n and j'_n are injections and the image of j'_n is $(W/N'_\chi W)^{(H_0, \tilde{\chi})}$.*

Let $\phi'_n : G \longrightarrow \mathbb{C}$ be defined by

$$\phi'_n(\gamma) = \begin{cases} \chi'_n(\gamma^{-1}), & \text{if } \gamma \in G'_n \\ 0, & \text{otherwise.} \end{cases}$$

Consider the corresponding function $\tilde{\phi}'_n : \tilde{G} \longrightarrow \mathbb{C}$.

Lemma 9. *Consider a pair (Y, φ) such that $O = O_Y$ is a maximal in $\mathcal{N}_{\mathrm{tr}}(\pi)$. Then for large enough n:*

(1) *Let $\mathcal{Y}_n \subset G'_{n+1} \cap G(Y)$ be a set of representatives for the G'_n double cosets in $G'_n(G_{n+1} \cap G(Y))G'_n$. Then large enough n,*

$$\tilde{\phi}'_{n+1} * \tilde{\phi}'_{n,n+1} * \tilde{\phi}'_n(g) = \begin{cases} \lambda \cdot (\chi'_n)^{-1}(h_1 h_2), & \text{if } g = h_1 y h_2 \text{ with } y \in \mathcal{Y}_n, h_1, h_2 \in G'_n \\ 0, & \text{if } g \notin G'_n \mathcal{Y}_n G'_n \end{cases}$$

where $\lambda = \operatorname{meas}(G'_n \cap G'_{n+1}) \operatorname{meas}(G'_n)$.

(2) *For large n, $I'_{n,n+1}$ is injective.*

Proof. From part (a) of Lemma 9 in [8], we have

$$\phi'_{n+1} * \phi'_{n,n+1} * \phi'_n(g) = \begin{cases} \lambda \cdot (\chi'_n)^{-1}(h_1 h_2), & \text{if } g = h_1 y h_2 \text{ with } y \in \mathcal{Y}_n, h_1, h_2 \in G'_n \\ 0, & \text{if } g \notin G'_n \mathcal{Y}_n G'_n \end{cases}$$

where $\lambda = \text{meas}(G'_n \cap G'_{n+1})\text{meas}(G'_n)$. Now part 1 follows from Lemma 4, as we have

$$\tilde{\phi}'_{n+1} * \tilde{\phi}'_{n,n+1} * \tilde{\phi}'_n = \widetilde{(\phi'_{n+1} * \phi'_{n,n+1} * \phi'_n)}. \tag{10}$$

Now we prove part 2. It is enough to say that $\pi(\tilde{\phi}'_{n+1} * \tilde{\phi}'_{n,n+1} * \tilde{\phi}'_n)$ acts by a non-zero multiple of identity on W'_n. This implies that $I'_{n+1,n} \circ I'_{n,n+1}$ is a non-zero multiple of identity on W'_n. From part 1 we get that $\tilde{\phi}'_{n+1} * \tilde{\phi}'_{n,n+1} * \tilde{\phi}'_n$ is a positive linear combination of functions $\tilde{\phi}'_{n,y} : \gamma \mapsto \tilde{\phi}'_n(\gamma y^{-1})$, where $y \in G_{n+1} \cap G(Y)$ is fixed and $G(Y)$ is centralizer of Y in G. Then the lemma follows from the fact that $\pi(y)$ acts trivially on W'_n for large n, so that

$$\pi(\tilde{\phi}'_{n,y})|_{W'_n} = \pi(\tilde{\phi}'_n)\pi(y)|_{W'_n} = \pi(\tilde{\phi}'_n)|_{W'_n}. \qquad \square$$

Theorem 3. *Let (π, W) be an irreducible admissible genuine representation of $\tilde{G}$.*

1. *The set of maximal elements in $\mathcal{N}_{\text{tr}}(\pi)$ coincides with the set of maximal elements in $\mathcal{N}_{Wh}(\pi)$.*
2. *Let $\mathcal{O}$ be a maximal element in $\mathcal{N}_{\text{tr}}(\pi)$. Then the coefficient $c_{\mathcal{O}}$ equals the dimension of the space of degenerate Whittaker forms with respect to any pair (Y, φ) such that with $Y \in \mathcal{O}$ is arbitrary and $\varphi : \mathbb{G}_m \longrightarrow \mathbf{G}$ satisfies $\text{Ad}(\varphi(s))Y = s^{-2}Y$ for all $s \in E^\times$.*

Proof. Let $\mathcal{O}$ be a maximal element in $\mathcal{N}_{\text{tr}}(\pi)$. Choose (Y, φ) such that $Y \in \mathcal{O}$ and $\varphi : \mathbb{G}_m \longrightarrow \mathbf{G}$ satisfying $\text{Ad}(\varphi(s))Y = s^{-2}Y$. Then, from Lemma 7, for large n we have

$$\dim W_n = c_{\mathcal{O}}.$$

Therefore $W_n \neq 0$ (resp $W'_n \neq 0$) for large n . From Lemma 8, the map j'_n is injective and maps surjectively onto $(W/N'_\chi W)^{(H_0,\tilde{Y})}$. But from second part of Lemma 9 and Equation 8, $I'_{n,m}$ is injective for large n and $m > n$ which implies that $W'_{n,\chi} = \cup_{m>n} \ker(I'_{n,m}) = 0$. Thus $\dim \mathcal{W} = \dim W'_n = \dim W_n = c_{\mathcal{O}}$, which proves part 2 of the theorem. In particular, $\mathcal{W} \neq 0$ and hence $\mathcal{O} \in \mathcal{N}_{\text{Wh}}(\pi)$. Now we claim that $\mathcal{O}$ is maximal in $\mathcal{N}_{\text{Wh}}(\pi)$. If not, there is a maximal orbit $\mathcal{O}' \in \mathcal{N}_{\text{Wh}}(\pi)$ such that $\mathcal{O} \lneq \mathcal{O}'$. From Proposition 1, there is $\mathcal{O}'' \in \mathcal{N}_{\text{tr}}(\pi)$ such that $\mathcal{O}' \leq \mathcal{O}''$. Therefore $\mathcal{O} \lneq \mathcal{O}''$ and $\mathcal{O}, \mathcal{O}'' \in \mathcal{N}_{\text{tr}}(\pi)$, a contradiction to the maximality of $\mathcal{O}$ in $\mathcal{N}_{\text{tr}}(\pi)$.

Let $\mathcal{O}$ be a maximal element in $\mathcal{N}_{\text{Wh}}(\pi)$. From Proposition 1, there is an element in $\mathcal{O}' \in \mathcal{N}_{\text{tr}}(\pi)$ such that $\mathcal{O} \leq \mathcal{O}'$. Take such a maximal $\mathcal{O}'$. Then by the argument in the above paragraph, $\mathcal{O}'$ is a maximal element in $\mathcal{N}_{\text{Wh}}(\pi)$. But $\mathcal{O}$ is also maximal in $\mathcal{N}_{\text{Wh}}(\pi)$. Hence $\mathcal{O} = \mathcal{O}'$. This proves that $\mathcal{O}$ is a maximal element in $\mathcal{N}_{\text{tr}}(\pi)$ too. $\square$

References

[1] W.-W. Li: *La formule des traces pour les revêtements de groupes réductifs connexes. II. Analyse harmonique locale* Ann. Scient. Éc. Norm. Sup. 4 e série, **45**, 787-859, (2012).

[2] W.-W. Li: *La formule des traces pour les revłtements de groupes réductifs connexes. I. Le développement géométrique fin*, Journal für die reine und angewandte Mathematik. **686**, 37-109, (2014).

[3] C. Mœglin, M. F. Vigneras and J.-L. Waldspurger: *Correspondances de Howe sur un corps p-adique*, Lecture notes in Mathematics, Springer-Verlag, Berlin, **1291**, (1987).

[4] C. Mœglin, and J.-L. Waldspurger: *Modèles de Whittaker dégénérés pour de groupes p-adiques*, Mathematics Zeitschrift, **196**, 427-452, (1987).
[5] C. Mœglin and J.-L. Waldspurger: *Spectral decomposition and Eisenstein series*, Appendix I, Lifting of Unipotent subgroups into a central extension, Cambridge tracts in Mathematics, Cambridge University press, Cambridge, **113**, 273-277, (1995).
[6] D. Prasad: *Invariant forms for the representations of GL_2 over a local field*. American Journal of Mathematics, **114**, 1317-1363, (1992).
[7] F. Rodier: *Modèles de Whittaker et caractéres de représentations*, Non commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1974), Lecture Notes in Mathematics, Springer, Berlin, **466** 151-171, (1975).
[8] Sandeep Varma: *On a result of Mœglin and Waldspurger in residual characteristic 2*, To appear in Mathematische Zeitschrift.

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Colaba, Mumbai - 400005, INDIA
E-mail address: shiv@math.tifr.res.in